CURRENT FDA-APPROVED MONO & DUAL AGONISTS*1-3,9

				FDA APPROVAL INDICATION	KEY STUDY		
	CLASS	GENERIC NAME	BRAND NAME	INDICATION (as of August 2023)^	SERIES NAME(S)+	DOSE RANGE/ FREQUENCY	DEVICE DESCRIPTION
	Mono GLP-1 agonists (expected Alc lowering: 0.5% to 1.6%)	exenatide (immediate release)	Byetta (injectable)	10 years old and older, type 2 diabetes, adjunct to diet and exercise#, glycemic control	EXSCEL	5 or 10 mcg twice/day, take within 60 min before eating	Pen (requires pen needles prescription)
		exenatide XR	Bydureon (injectable)	10 years old and older, type 2 diabetes, adjunct to diet and exercise#, glycemic control	DURATION	2 mg weekly	Pen injector-Bydureon BCise
		liraglutide	Victoza (injectable)	10 years old and older, type 2 diabetes, adjunct to diet and exercise#, glycemic control, reduces risk of major adverse cardiovascular events and established CVD	LEADER	0.6, 1.2, or 1.8 mg daily (doesn't have to be taken before meals)	Pen (requires pen needles prescription)
		liraglutide	Saxenda (injectable)	Adults: BMI > 30 or > 27 if weight-related medical issues Pediatric: 12 years old and older, initial weight > 60 kg (132 lbs)	SCALE	3 mg maximum dose daily	Pen (requires pen needles prescription)
i J		dulaglutide	Trulicity (injectable)	Adults, type 2 diabetes, adjunct to diet and exercise#, glycemic control, reduces risk of major adverse cardiovascular events and established CVD, or multiple risk factors	REWIND	0.75, 1.5, 3, or 4.5 mg weekly	Pen injector
		semaglutide	Ozempic (injectable)	Adults, type 2 diabetes, adjunct to diet and exercise#, glycemic control, reduces risk of major adverse cardiovascular events and established CVD	SUSTAIN, SELECT	0.25, 0.5, 1, or 2 mg weekly	Pen (comes with needles)
		semaglutide	Rybelsus (oral)	Adults, type 2 diabetes, adjunct to diet and exercise#, glycemic control	PIONEER	3, 7, or 14 mg daily in the morning. Take on empty stomach alone with 4 oz water, at least 30 minutes before eating and drinking	N/A (an oral medication)
		semaglutide	Wegovy (injectable)	Adults: BMI > 30 or > 27 if weight-related medical issues. Pediatric: 12 years old and older, initial BMI 95 th percentile, age, sex	STEP	2.4 mg maximum dose	Pen (comes with hidden needle on the pen)
	Dual GLP-1 agonist/ GIP (expected A1c lowering: 1.8% to 2.4%)	tirzepatide	Mounjaro (injectable)	Type 2 diabetes	SURPASS, SURMOUNT	2.5, 5, 7.5, 10, 12.5, or 15 mg daily	Pen injector

*LISTED CHRONOLOGICALLY AS THEY WERE APPROVED BY THE FDA.

[•]Details on FDA approval indication(s) as of August 2023. With filed research outcomes, the FDA can approve additional indications that are then integrated into the package insert. The FDA can issue new label warnings for FDA-approved medications at any time. In late September 2023, the FDA added a warning on the label of semaglutide (Ozempic), telling users of the potential for an increased risk of lieus (an intestinal blockage). The FDA added a warning on the label of semaglutide (Ozempic), telling users of the potential for an increased risk of lieus (an intestinal blockage). The FDA added a warning so the babel of semaglutide (Ozempic), telling users of the potential for an increased risk of lieus (an intestinal blockage). The FDA added a warning so the tota warning label for Mourjaro and Wegovy. Reference: Drug safety-related labeling changes (SRLC). FDA website. https://www.accessdata.fda.gov/scripts/cder/safetylabelingchanges/index.cfm?event-searchdetail.page&DrugNameID=2183. Updated September 22, 2023. Accessed October 4, 2023.

#The term "diet and exercise" has been used by the FDA for medication approvals for decades.

+These acronyms for studies represent some, but not all, of the series of studies conducted in the research and development for each medication to support FDA approval and use indications and approval by other global medication institutions.